

Research

Outcomes of conversion total hip arthroplasty following previous hip fracture surgery
Bone Joint Open 2025 Feb 14;6(2):195-205.

The incidence and risk factors for abnormal postoperative blood tests following primary total joint replacement
Bone Joint Open 2023 Nov 24;4(11):899-905.

Clinical and cost effectiveness of single stage compared with two stage revision for hip prosthetic joint infection (INFORM): pragmatic, parallel group, open label, randomised controlled trial
BMJ 2022 Oct 31:379.

The STAR care pathway for patients with pain at 3 months after total knee replacement: a multicentre, pragmatic, randomised, controlled trial
Lancet Rheumatol. 2022 Jan 28;4(3):e188-e197.

Greater trochanteric pain syndrome: focused shockwave therapy versus an ultrasound guided injection: a randomised control trial
Hip Int 2023 May;33(3):490-499.

Elective surgery cancellations due to the COVID-19 pandemic: global predictive modelling to inform surgical recovery plans. COVID Surg Collaborative.
Br J Surg. 2020 May 12:10.1002.

Prospective observational cohort study on grading the severity of postoperative complications in global surgery research. International Surgical Outcomes Study (ISOS) group.
Br J Surg. 2019 Jan;106(2):e73-e80.

Risk factors associated with revision for prosthetic joint infection after hip replacement: a prospective observational cohort study.
Lancet Infect Dis. 2018 Sep;18(9):1004-1014.

One- and two-stage surgical revision of peri-prosthetic joint infection of the hip: a pooled individual participant data analysis of 44 cohort studies.
Eur J Epidemiol. 2018 Oct;33(10):933-946.

The surgical safety checklist and patient outcomes after surgery: a prospective observational cohort study, systematic review and meta-analysis. International Surgical Outcomes Study (ISOS) group.
Br J Anaesth. 2018 Jan;120(1):146-155.

Critical care admission following elective surgery was not associated with survival benefit: prospective analysis of data from 27 countries. Surgical Outcomes Study (ISOS) group.
Intensive Care Med. 2017 Jul;43(7):971-979.

Global patient outcomes after elective surgery: prospective cohort study in 27 low-, middle- and high-income countries. International Surgical Outcomes Study group.
Br J Anaesth. 2016 Oct 31;117(5):601-609.

Re-Infection Outcomes Following One- And Two-Stage Surgical Revision of Infected Knee Prosthesis: A Systematic Review and Meta- Analysis
PLoS ONE. 2016 11(3): e0151537.

Patient-Related Risk Factors for Periprosthetic Joint Infection after Total Joint Arthroplasty: A Systematic Review and Meta-Analysis
PLoS ONE. 2016 11(3): e0150866.

One-stage or two-stage revision surgery for prosthetic hip joint infection--the INFORM trial: a study protocol for a randomised controlled trial.
Trials. 2016 Feb 17;17:90.

Re-Infection Outcomes following One- and Two-Stage Surgical Revision of Infected Hip Prosthesis: A Systematic Review and Meta- Analysis
PLoS ONE. 2015 10(9): e0139166.

Re-infection outcomes following one- and two-stage surgical revision of infected hip prosthesis in unselected patients: protocol for a systematic review and an individual participant data meta-analysis.
Syst Rev. 2015 Apr 25;4:58.

Human osteoblast growth and maturation in response to metformin and the ienopyridone A769662
SAJ Pharmacy and Pharmacology 2015, 1:1.

High failure rates of the Spectron EF stem at a minimum of 10 year's follow up.
J Arthroplasty. 2014 Oct;29(10):1956-60.

Mid-term review of ADEPT metal-on-metal hip prosthesis. Functional, radiological and metal ion analysis
Open Journal of Orthopaedics. 2014; 4: 38-43.

Clinical and radiological results of the collarless polished tapered stem at 15 years follow-up.
J Bone Joint Surg Br. 2012 Jul;94(7):889-94.

Radiographic landmarks for tunnel placement in reconstruction of the medial patellofemoral ligament.
Knee Surg Sports Traumatol Arthrosc. 2012 Dec;20(12):2380-4.

Cemented polished tapered hip replacements in patients aged 50 years or less. A minimum 10-year follow up.
J Arthroplasty. 2010; 25: 692-9.

Large diameter femoral head uncemented total hip replacement to treat fractured neck of femur.
Injury. 2009; 40: 752-5.

The economic implications of transfusion post primary total knee arthroplasty – an argument for autologous blood retransfusion?
JBJS (Br). 2009; 91B: Supp II, 356.

Collarless Polished Tapered Stem (CPT). Clinical and Radiological Follow-up at a Minimum of 10 Years.
J Bone Joint Surg (Br). 2008; 90B: 16-22.

Cement burn of the skin during hip arthroplasty.
Ann R Coll Surg Engl. 2007; 89: 151-152.

The Exeter Universal stem. A minimum 10-year review from an independent centre.
J Bone Joint Surg (Br). 2006; 88B: 1584-1590.

Failure of surgery for scaphoid non-union is associated with smoking.
J Hand Surg (Br). 2006; 31B: 252-255.

Gradenigo's syndrome: successful conservative treatment in adult and paediatric patients.
J Laryngol Otol. 2005; 119: 325-329.

Genetic influences in the aetiology of tears of the rotator cuff.
J Bone Joint Surg (Br). 2004; 86B: 696-700.

Determination of a standard site for the measurement of bone mineral density of the human calcaneus.
J Anat. 1998; 193: 449-456.